

Infants', Children's and Jr. TYLENOL®

Dosing Information for Healthcare Professionals



Use this chart to determine the proper dose of TYLENOL® for your patient. If possible, use weight to dose; otherwise use age.

DOSE — Every 4 hours as needed. Do not give more than 5 doses in 24 hours.

		Infants' TYLENOL® Oral Suspension	Children's TYLENOL® Oral Suspension	Children's TYLENOL® MELTAWAYS® Chewable Tablets	Jr. TYLENOL® MELTAWAYS® Chewable Tablets
		Active Ingredient: Acetaminophen 160 mg (in each 5mL or 1 tsp)	Active Ingredient: Acetaminophen 160 mg (in each 5mL or 1 tsp)	Active Ingredient: Acetaminophen 80 mg (in each tablet)	Active Ingredient: Acetaminophen 160 mg (in each tablet)
Weight (lbs)	Age				
6-11 lbs	0-3 mos	1.25 mL	—	—	—
12-17 lbs	4-11 mos	2.5 mL	—	—	—
18-23 lbs	12-23 mos	3.75 mL	—	—	—
24-35 lbs	2-3 yrs	5 mL	5 mL (1 tsp)	2 tablets	—
36-47 lbs	4-5 yrs	—	7.5 mL (1½ tsp)	3 tablets	—
48-59 lbs	6-8 yrs	—	10 mL (2 tsp)	4 tablets	2 tablets
60-71 lbs	9-10 yrs	—	12.5 mL (2½ tsp)	5 tablets	2½ tablets
72-95 lbs	11 yrs	—	15 mL (3 tsp)	6 tablets	3 tablets

mL = milliliter tsp = teaspoonful

See reverse side for important information ▶▶

Remind parents and caregivers to:

- Read and follow the label on all **TYLENOL®** products
- Take every 4 hours as needed. Do **NOT** exceed more than 5 doses in 24 hours
- Do **NOT** use with any other product containing acetaminophen
- Use only the dosing device that comes with a specific product
- **Children's TYLENOL® MELTAWAYS® Chewable Tablets (80 mg/tablet) are not the same concentration as Jr. TYLENOL® MELTAWAYS® Chewable Tablets (160 mg/tablet)**
- **All Infants' and Children's TYLENOL® Oral Suspension products have the same acetaminophen concentration (160 mg/5 mL)**

For more information, visit TylenolProfessional.com

Children's
TYLENOL